

# MPOX, AIDS, COVID-19 SHOW THE CHALLENGES OF PUBLIC HEALTH MESSAGING

Relevant for: Developmental Issues | Topic: Health & Sanitation and related issues

To enjoy additional benefits

CONNECT WITH US

December 23, 2022 05:56 pm | Updated 05:56 pm IST

COMMENTS

SHARE

READ LATER

Inclusive public health messaging can motivate the public to make better decisions regarding their personal health and the health of others. Image for Representation. | Photo Credit: Gopal Nagara

During infectious disease outbreaks, clinicians and public health officials are tasked with providing accurate guidance for the public on how to stay safe and protect themselves and their loved ones. However, sensationalised media coverage can distort how the public perceives new emerging infections, including where they come from and how they spread. This can foster fear and stigma, especially toward communities that are already mistrustful of the health care system.

The racial and sexual stigma surrounding monkeypox is what spurred the World Health Organization (WHO) to [rename](#) the disease to mpox in November 2022. While this is a step in the right direction, I believe more work needs to be done to reduce the stigma surrounding infectious diseases like mpox.

I am an infectious disease researcher who studies HIV, COVID-19 and mpox. During the COVID-19 pandemic, I was the lead investigator at the University of Pittsburgh for a national survey looking at how COVID-19 has affected different communities. Effective public health communication isn't easy when conflicting messages may come from many sources, including family and friends, other community members or the internet. But there are ways that public health officials can make their own messaging more inclusive while mitigating stigma.

Inclusive public health messaging can motivate the public to make better decisions regarding their personal health and the health of others. This effort often involves engaging the communities most affected by an outbreak. Unfortunately, because these communities are heavily affected by the infection and tend to experience some form of inequity, they are often blamed by society for spreading the disease.

COVID-19 drove an increase in hate crimes related to the pandemic against Chinese and other Asian communities in the United States. A 2022 UCLA survey found that 8% of Asian American and Pacific Islander adults in California experienced a COVID-19 related hate incident.

Effective public health messaging can focus on the fact that while infections may first affect

certain groups of people, they often spread to other groups and eventually encompass entire communities. Infections are caused by bacteria, viruses and fungi. They don't discriminate by race, gender or sexual orientation. Messages that focus on the pathogens, rather than the communities, may reduce stigma.

[Also Read | Omicron subvariant BF.7, linked to China spike, first found in India in July](#)

Visually inclusive messages are also likely to engage a greater portion of the community. Examples include making sure that the people represented in posters and flyers, images on TV and websites, and other informational material are from diverse backgrounds. This sends a more unified message that what affects an individual also affects the larger community.

Many media outlets, especially on social media, use fear-based messaging to report on infectious diseases. While this may reinforce certain protective behaviours, such as using condoms during sex, it may also increase stress and anxiety. Fear-based messages also worsen stigma, leading to increased discrimination against communities that are already vulnerable and mistrustful of health care. Ultimately, this leads people to avoid seeking health care and can worsen health outcomes.

Public health officials have often used fear-based messaging in response to sexually transmitted infections, or STIs, like HIV, chlamydia and gonorrhoea. Sex itself is highly stigmatised by society. I have found that some of my patients would prefer to avoid getting tested and treated for an STI rather than deal with the shame of having an STI.

Making sexual health and STI testing routine and integral parts of overall wellness and health is an important step to reduce the stigma around them. Similarly, messaging that normalises the challenges faced by people at risk for certain infections could help avoid causing shame.

Infections affect different people differently. COVID-19 might be a mildly stuffy nose for one person, and it could be months in an intensive care unit hooked up to a ventilator for another. Messages that focus on the successes of medical and public health interventions that resonate with communities are most likely to be successful.

Different groups have different exposure risks as well. Mpox heavily affected gay and bisexual men in 2022. One reason why was related to how the virus is transmitted. Prior research suggested that mpox was largely transmitted by close skin-to-skin contact, but emerging studies raised the question of whether the 2022 outbreaks were being driven more by sexual transmission.

[Also Read | HIV drugs shortage is a challenge to ending AIDS in India](#)

There was controversy as to whether public health messaging should highlight sexual encounters as a potential transmission route. This can risk further stigmatising gay and bisexual men versus potentially overlooking these key at-risk populations. Some advocates argued that promoting the message that mpox was primarily transmitted by close contact would prevent resources and interventions from reaching the groups of people most affected by the disease.

One size does not always fit all when it comes to public health messaging. Multiple messages may be necessary for different groups of people based on their risk of infection or severe disease. An August 2022 Centers for Disease Control and Infection survey found that 50% of gay and bisexual men reduced their sexual encounters in response to the mpox outbreak. Since late summer, mpox rates have been dropping rapidly, and many experts think that both behaviour change and vaccination may have contributed to the falling rates. Studies like these

further support the importance of directly engaging with communities to encourage healthy behaviour change.

Mistrust is also a barrier to effective messaging. Some communities may be mistrustful of medical and health care systems because of prior histories of exploitation, such as the Tuskegee study, where researchers prevented Black participants from receiving syphilis treatment for decades in the mid-20th century, and ongoing fear of mistreatment.

Identifying trusted community champions and health care providers – especially ones who belong to that community – to deliver a public health message may increase its acceptance. One 2019 study, for example, found that Black men were more likely to accept vaccines, medical advice and engage in health care services if they had a Black health care provider.

Effectively delivering public health messaging is a complicated and challenging process. But talking to and listening to the communities most affected by an outbreak can make a difference.

*Ken Ho, University of Pittsburgh Health Sciences (The Conversation)*

COMMENTS

SHARE

[Coronavirus](#) / [Monkeypox](#) / [AIDS](#) / [health](#) / [discrimination](#)

BACK TO TOP

Comments have to be in English, and in full sentences. They cannot be abusive or personal. Please abide by our [community guidelines](#) for posting your comments.

We have migrated to a new commenting platform. If you are already a registered user of The Hindu and logged in, you may continue to engage with our articles. If you do not have an account please register and login to post comments. Users can access their older comments by logging into their accounts on Vuukle.

**END**

Downloaded from [crackIAS.com](#)

© **Zuccess App** by crackIAS.com

